

BIRTH DEFECT RISK FACTOR SERIES:

HYDROCEPHALY

DEFINITION

Congenital hydrocephaly is one of the most common central nervous system anomalies. Hydrocephaly is an enlargement of the head caused by an abnormal accumulation of cerebrospinal fluid (CSF) in the cranium due to an imbalance between the production and absorption of CSF. This forces the ventricles to enlarge (ventricular dilatation or ventriculomegaly), which in turn exerts pressure on the surrounding brain tissue, causing the brain tissue to shrink and the head to enlarge (Schrandt-Stumpel and Fryns, 1998; Buyse, 1990; Vintzileos et al., 1983). Congenital hydrocephaly usually develops by the twentieth week of gestation (Stoll et al., 1992a).

There are various types or classifications of congenital hydrocephaly. Aqueductal stenosis is a type of hydrocephaly which results from narrowing of the aqueduct of Sylvius, an opening connecting the third and fourth ventricles in the brain. It is the most common form of hydrocephaly. Dandy-Walker syndrome is a group of defects consisting of enlargement of the fourth ventricle of the brain, complete or partial absence of the cerebellar vermis (the middle area between the two cerebral hemispheres), cyst of the posterior fossa (internal base of the skull), and hydrocephaly. The hydrocephaly associated with Dandy-Walker syndrome may not be present at birth but develop later (Buyse, 1990). Dandy-Walker syndrome accounts for 5-12% of hydrocephaly (Vintzileos et al., 1983).

Congenital hydrocephaly is heterogeneous in origin. The defect can be associated with chromosomal abnormalities (trisomy 21, trisomy 13, trisomy 18, triploidy, etc.) and other syndromes (Walker-Warburg syndrome, Meckel syndrome, Smith-Lemli-Opitz syndrome, and chondrodystrophies, etc.). Some cases seem to be linked to the X chromosome (Torfs and Christianson, 1998). About one-quarter of the infants with hydrocephaly also have spina bifida; conversely, approximately 80% of children with spina bifida also have hydrocephaly. Hydrocephaly can also be secondary to central nervous system anomalies (encephalocele, holoprosencephaly, neoplasms, etc.). Most congenital hydrocephaly cases have additional birth defects such as congenital heart disease and cleft lip and/or palate (Schrandt-Stumpel and Fryns, 1998; Stoll et al., 1992a; Buyse, 1990; Drugan et al., 1989; Hudgins et al., 1988; Vintzileos et al., 1983; Chervenak et al., 1985).

Hydrocephaly (ventriculomegaly) can be detected prenatally by ultrasonography (Fadel, 1989; Vintzileos et al., 1987). Studies from various birth defects surveillance systems have found that, in regions where elective termination is allowed, prenatal diagnosis and elective termination reduce the birth prevalence of congenital hydrocephaly (Riley et al., 1998; Julian-Reynier et al., 1994; Stoll et al., 1995; Stoll et al., 1992a; Stoll et al., 1992b; Stone et al., 1989; Drugan et al., 1989; Hudgins et al., 1988; Chervenak et al., 1985).

DEMOGRAPHIC AND REPRODUCTIVE FACTORS

One study reported **racial/ethnic** differences in hydrocephaly rates, with the rate being highest for Native Americans, followed by African-Americans, whites, Asians, and Hispanics (Chavez et al., 1988). Another study found hydrocephaly risk to be lower among infants of Vietnamese mothers when compared with infants of non-Hispanic white mothers (Shaw et al., 2002). However, other investigations did not identify any racial/ethnic differences in hydrocephaly risk (Leck and Lancashire, 1995; Stoll et al., 1992a; Wiswell et al., 1990; Finley et al., 1980).

A **secular trend** for congenital hydrocephaly has been reported by one study, where a decline in prevalence over time was found, paralleling the decline in neural tube defect prevalence (Stone et al.,

1989). However, another investigation observed no trend (Wiswell et al., 1990). There does not appear to be any **seasonal variation** in congenital hydrocephaly prevalence (Stoll et al., 1992a; Castilla et al., 1990; Wiswell et al., 1990; Bound et al., 1989).

Congenital hydrocephaly prevalence does not appear to be influenced by **geographic location** (Stoll et al., 1992a). However, one study found a reduction in hydrocephaly risk with **higher altitudes** (Castilla et al., 1999).

Parental age does not clearly seem to affect risk of having a child with hydrocephaly (Hollier et al., 2000; McIntosh et al., 1995; Stoll et al., 1992a).

Information on the relationship between **infant sex** and hydrocephaly is mixed. Some investigations observed no association (Riley et al., 1998; Stoll et al., 1992a; Buyse, 1990), while other studies reported a predominance among males (Lary and Paulozzi, 2001; Wiswell et al., 1990; Lowry et al., 1986).

Investigations into the relationship between **plurality** and congenital hydrocephaly have produced inconsistent results (Riley et al., 1998; Stoll et al., 1992a; Doyle et al., 1991; Ramos-Arroyo, 1991; Kallen, 1986). Hydrocephaly has been associated with lower **birth weight** and **intrauterine growth retardation** (Riley et al., 1998; Mili et al., 1991; Khoury et al., 1988). One investigation reported no statistically significant relationship between hydrocephaly and **macrosomia** (Waller et al., 2001), while another study found increased risk of hydrocephaly with large for gestational age (Lapunzina et al., 2002).

A woman who has had one child with congenital hydrocephaly has a **recurrence risk** of 1-5% of having another affected child. If the hydrocephaly is associated with an inherited disorder, the risk is higher (Schrandt-Stumpel and Fryns, 1998; Buyse, 1990). Hydrocephaly rates have been reported to be higher for **consanguineous parents** (Rittler et al., 2001; Rajab et al., 1998; Stoltenberg et al., 1997; Stoll et al., 1992a).

FACTORS IN LIFESTYLE OR ENVIRONMENT

One investigation identified no significant association between **maternal nursing occupation** and hydrocephaly risk (Matte et al., 1993). Another study failed to identify any significant association between hydrocephaly and proximity to various types of **industry** (Castilla et al., 2000). In one investigation, risk of hydrocephaly was significantly increased with high environmental **lead** exposure (Vincenti et al., 2001).

Maternal prenatal **infection** (toxoplasmosis, syphilis, cytomegalovirus, rubella) has been strongly associated with increased hydrocephaly prevalence (Schrandt-Stumpel and Fryns, 1998; Stoll et al., 1992a; Vintzileos et al., 1983). One investigation failed to find any effect on hydrocephaly risk for maternal **epilepsy, x-rays, hypertension, fever, "flu," medication exposure, or occupational exposure** (Stoll et al., 1992a). The relationship between maternal **diabetes** and hydrocephaly risk is not clear (Stoll et al., 1992a; Becerra et al., 1990). Another study found no relationship between maternal **hypothyroidism** and hydrocephaly; however, risk of the defect was increased with maternal **hyperthyroidism** (Khoury et al., 1989). Maternal **common cold** in the first trimester of pregnancy has been reported to increase risk of hydrocephaly (Zhang and Cai, 1993).

Maternal **smoking** does not appear to affect hydrocephaly rates (Van Den Eeden et al., 1990). An investigation reported increased risk of hydrocephaly with maternal **oral contraceptive** use, although exposure was reported for only one case (Correy et al., 1991). Another study has found no relationship between **parental occupation of farmer** and hydrocephaly prevalence; however, the study suggested increased hydrocephaly risk with parental **pesticide** exposure (Kristensen et al., 1997). A recent study

reported an increased rate of hydrocephaly among infants born to women with **obesity** but who were not diabetic (Moore et al., 2000). Living in proximity to **hazardous waste sites** has not been found to affect risk of hydrocephaly (Dolk et al., 1998) nor has **paternal ionizing radiation exposure** (Doyle et al., 2000). One study reported no association between hydrocephaly risk and maternal or paternal occupational exposure to **electromagnetic fields**; however, exposure was based on linkage to census data and exposure assessments by an expert panel (Blaasaas et al., 2002). Hydrocephaly does not appear related to **water chlorination** (Kallen and Robert, 2000).

Case reports and a case-control study suggest that risk of hydrocephaly may be increased with maternal use of **misoprostol**, a synthetic prostaglandin used for elective terminations (Orioli and Castilla, 2000). Hydrocephaly has not been associated with maternal use of cough medicines containing **dextromethorphan** (Martinez-Frias and Rodriguez-Pinilla, 2001) the antibiotic **oxytetracycline** (Czeizel and Rockenbauer, 2000), **cephalosporin antibiotics** (Czeizel et al., 2001a), **nalidixic acid** (Czeizel et al., 2001b), ampicillin (Czeizel et al., 2001c), **calcium channel blockers** (Sorensen et al., 2001), or the **benzodiazepines** nitrazepam, medazepam, tofisopam, alprazolam, and clonazepam (Eros et al., 2002). One investigation identified a relationship between maternal **anesthesia** exposure in the first trimester and hydrocephaly, particularly in association with other birth defects (Sylvester et al., 1994).

Several studies have evaluated the relationship between maternal **multivitamin use** and **folic acid** and congenital hydrocephaly risk. The studies found a slight reduction in risk; however, this reduction was not considered significant (Werler et al., 1999; Czeizel et al., 1996; Czeizel, 1993). Furthermore, a study that examined **co-trimoxazole**, a combination of trimethoprim and sulfamethoxazole that is a folic acid antagonist, failed to find any association between the medication and hydrocephaly (Czeizel, 1990).

REFERENCES

-
- | | |
|--|---|
| <p>Becerra JE, Khoury MJ, Cordero JF, Erickson JD. Diabetes mellitus during pregnancy and the risks for specific birth defects: a population-based case-control study. <i>Pediatrics</i> 1990;85:1-9.</p> <p>Blaasaas KG, Tynes T, Irgens A, Lie RT. Risk of birth defects by parental occupational exposure to 50 Hz electromagnetic fields: a population-based study. <i>Occup Environ Med</i> 2002;59:92-97.</p> <p>Bound JP, Harvey PW, Francis BJ. Seasonal prevalence of major congenital malformations in the Fylde of Lancashire 1957-1981. <i>J Epidemiol Community Health</i> 1989;43:330-342.</p> <p>Buyse ML, editor-in-chief. <i>Birth Defect Encyclopedia</i>. Cambridge, Massachusetts: Blackwell Scientific Publications, 1990.</p> <p>Castilla EE, Campana H, Camelo JS. Economic activity and congenital anomalies: an ecologic study in Argentina. <i>Environ Health Perspect</i> 2000;108:193-197.</p> | <p>Castilla EE, Lopez-Camelo JS, Campana H. Altitude as a risk factor for congenital anomalies. <i>Am J Med Genet</i> 1999;86:9-14.</p> <p>Castilla EE, Orioli IM, Lugarinho R, Dutra GP, Lopez-Camelo JS, Campana HE, Spagnolo A, Mastroiacovo P. Monthly and seasonal variations in the frequency of congenital anomalies. <i>Int J Epidemiol</i> 1990;19:399-404.</p> <p>Chavez GF, Cordero JF, Becerra JE. Leading major congenital malformations among minority groups in the United States, 1981-1986. <i>Mor Mortal Wkly Rep CDC Surveill Summ</i> 1988;37:17-24.</p> <p>Chervenak FA, Berkowitz RL, Tortora M, Hobbins JC. The management of fetal hydrocephalus. <i>Am J Obstet Gynecol</i> 1985;151:933-942.</p> <p>Correy JF, Newman NM, Collins JA, Burrows EA, Burrows RF, Curran JT. Use of prescription drugs in the first trimester and congenital malformations. <i>Aust N Z J Obstet Gynaecol</i></p> |
|--|---|

1991;31:340-344.

Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. Use of cephalosporins during pregnancy and in the presence of congenital abnormalities: a population-based, case-control study. *Am J Obstet Gynecol* 2001a;184:1289-1296.

Czeizel AE, Sorensen HT, Rockenbauer M, Olsen J. A population-based case-control teratologic study of nalidixic acid. *Int J Gynaecol Obstet* 2001b;73:221-228.

Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. A population-based case-control teratologic study of ampicillin treatment during pregnancy. *Am J Obstet Gynecol* 2001c;185:140-147.

Czeizel AE, Rockenbauer M. A population-based case-control teratologic study of oral oxytetracycline treatment during pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2000;88:27-33.

Czeizel AE, Toth M, Rockenbauer M. Population-based case control study of folic acid supplementation during pregnancy. *Teratology* 1996;53:345-351.

Czeizel AE. Prevention of congenital abnormalities by periconceptional multivitamin supplementation. *BMJ* 1993;306:1645-1648.

Czeizel A. A case-control analysis of the teratogenic effects of co-trimoxazole. *Reprod Toxicol* 1990;4:305-313.

Dolk H, Vrijheid M, Armstrong B, Abramsky L, Bianchi F, Garne E, Nelen V, Robert E, Scott JE, Stone D, Tenconi R. Risk of congenital anomalies near hazardous-waste landfill sites in Europe: the EUROHAZCON study. *Lancet* 1998;352:423-427.

Doyle P, Maconochie N, Roman E, Davies G, Smith PG, Beral V. Fetal death and congenital malformation in babies born to nuclear industry employees: report from the nuclear industry family study. *Lancet* 2000;356:1293-1299.

Doyle PE, Beral V, Botting B, Wale CJ. Congenital malformations in twins in England and Wales. *J Epidemiol Community Health* 1991;45:43-48.

Drugan A, Krause B, Canady A, Zador IE, Sacks AJ, Evans MI. The natural history of prenatally diagnosed cerebral ventriculomegaly. *JAMA* 1989;261:1785-1788.

Eros E, Czeizel AE, Rockenbauer M, Sorensen HT, Olsen J. A population-based case-control teratologic study of nitrazepam, medazepam, tofisopam, alprazolam and clonazepam treatment during pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2002;101:147-154.

Fadel HE. Antenatal diagnosis of fetal intracranial anomalies. *J Child Neurol* 1989;4:S107-S112.

Finley WH, Gustavson KH, Hall TM, Hurst DC, Barganier CM, Wiedmeyer JA. Birth defects surveillance: Jefferson County, Alabama, and Uppsala County, Sweden. *South Med J* 1994;87:440-445.

Hollier LM, Leveno KJ, Kelly MA, McIntire DD, Cunningham FG. Maternal age and malformations in singleton births. *Obstet Gynecol* 2000;96:701-706.

Hudgins RJ, Edwards MS, Goldstein R, Callen PW, Harrison MR, Filly RA, Golbus MS. Natural history of fetal ventriculomegaly. *Pediatrics* 1988;82:692-697.

Julian-Reynier C, Philip N, Scheiner C, Aurrant Y, Chabal F, Maron A, Gombert A, Ayme S. Impact of prenatal diagnosis by ultrasound on the prevalence of congenital anomalies at birth in southern France. *J Epidemiol Community Health* 1994;48:290-296.

Kallen B. Congenital malformations in twins: a population study. *Acta Genet Med Gemellol (Roma)* 1986;35:167-178.

Kallen BA, Robert E. Drinking water chlorination and delivery outcome-a registry-based study in Sweden. *Reprod Toxicol* 2000;14:303-309.

- Khoury MJ, Becerra JE, d'Almada PJ. Maternal thyroid disease and risk of birth defects in offspring: a population-based case-control study. *Paediatr Perinat Epidemiol* 1989;3:402-420.
- Khoury MJ, Erickson JD, Cordero JF, McCarthy BJ. Congenital malformations and intrauterine growth retardation: a population study. *Pediatrics* 1988;82:83-90.
- Kristensen P, Irgens LM, Andersen A, Bye AS, Sundheim L. Birth defects among offspring of Norwegian farmers, 1967-1991. *Epidemiology* 1997;8:537-544
- Lapunzina P, Lopez Camelo JS, Rittler M, Castilla EE. Risks of congenital anomalies in large for gestational age infants. *J Pediatr* 2002;140:200-204.
- Lary JM, Paulozzi LJ. Sex differences in the prevalence of human birth defects: a population-based study. *Teratology* 2001;64:237-251.
- Leck I, Lancashire RJ. Birth prevalence of malformations in members of different ethnic groups and in the offspring of matings between them, in Birmingham, England. *J Epidemiol Community Health* 1995;49:171-179.
- Lowry RB, Thunem NY, Silver M. Congenital anomalies in American Indians of British Columbia. *Genet Epidemiol* 1986;3:455-467.
- Martinez-Frias ML, Rodriguez-Pinilla E. Epidemiologic analysis of prenatal exposure to cough medicines containing dextromethorphan: No evidence of human teratogenicity. *Teratology* 2001;63:38-41.
- Matte TD, Mulinare J, Erickson JD. Case-control study of congenital defects and parental employment in health care. *Am J Ind Med* 1993;24:11-23.
- McIntosh GC, Olshan AF, Baird PA. Paternal age and the risk of birth defects in offspring. *Epidemiology* 1995;6:282-288.
- Mili F, Edmonds LD, Khoury MJ, McClearn AB. Prevalence of birth defects among low-birth-weight infants. A population study. *Am J Dis Child* 1991;145:1313-1318.
- Moore LL, Singer MR, Bradlee ML, Rothman KJ, Milunsky A. A prospective study of the risk of congenital defects associated with maternal obesity and diabetes mellitus. *Epidemiology* 2000;11:689-694.
- Orioli IM, Castilla EE. Epidemiological assessment of misoprostol teratogenicity. *BJOG* 2000;107:519-523.
- Rajab A, Vaishnav A, Freeman NV, Patton MA. Neural tube defects and congenital hydrocephalus in the Sultanate of Oman. *J Trop Pediatr* 1998;44:300-303
- Ramos-Arroyo MA. Birth defects in twins: study in a Spanish population. *Acta Genet Med Gemellol (Roma)* 1991;40:337-344.
- Riley MM, Halliday JL, Lumley JM. Congenital malformations in Victoria, Australia, 1983-95: an overview of infant characteristics. *J Paediatr Child Health* 1998;34:233-240.
- Rittler M, Liascovich R, Lopez-Camelo J, Castilla EE. Parental consanguinity in specific types of congenital anomalies. *Am J Med Genet* 2001;102:36-43.
- Schrander-Stumpel C, Fryns JP. Congenital hydrocephalus: nosology and guidelines for clinical approach and genetic counseling. *Eur J Pediatr* 1998;157:355-362.
- Shaw GM, Carmichael SL, Nelson V. Congenital malformations in offspring of Vietnamese women in California, 1985-97. *Teratology* 2002;65:121-124.
- Sorensen HT, Czeizel AE, Rockenbauer M, Steffensen FH, Olsen J. The risk of limb deficiencies and other congenital abnormalities in children exposed in utero to calcium channel blockers. *Acta Obstet Gynecol Scand* 2001;80:397-401.
- Stoll C, Dott B, Alembik Y, Roth MP. Evaluation of routine prenatal diagnosis by a registry of

congenital anomalies. Prenat Diagn 1995;15:791-800.

Stoll C, Alembik Y, Dott B, Roth MP. An epidemiologic study of environmental and genetic factors in congenital hydrocephalus. Eur J Epidemiol 1992a;8:797-803.

Stoll C, Alembik Y, Dott B, Roth MP, Finck S. Evaluation of prenatal diagnosis by a registry of congenital anomalies. Prenat Diagn 1992b;12:263-270.

Stoltenberg C, Magnus P, Lie RT, Daltveit AK, Irgens LM. Birth defects and parental consanguinity in Norway. Am J Epidemiol 1997;145:439-448.

Stone DH, Womersley J, Sinclair T, Evans L. Declining prevalence of hydrocephalus. Eur J Epidemiol 1989;5:398-399.

Sylvester GC, Khoury MJ, Lu X, Erickson JD. First-trimester anesthesia exposure and the risk of central nervous system defects: a population-based case-control study. Am J Public Health 1994;84:1757-1760.

Torfs CP, Christianson RE. Anomalies in Down syndrome individuals in a large population-based registry. Am J Med Genet 1998;77:431-438.

Van Den Eeden, SK, Karagas MR, Daling JR,

Vaughan TL. A case-control study of maternal smoking and congenital malformations. Paediatr Perinat Epidemiol 1990;4:147-155.

Vinceti M, Rovesti S, Bergomi M, Calzolari E, Candela S, Campagna A, Milan M, Vivoli G. Risk of birth defects in a population exposed to environmental lead pollution. Sci Total Environ 2001;278:23-30.

Vintzileos AM, Ingardia CJ, Nochimson DJ. Congenital hydrocephalus: A review and protocol for perinatal management. Obstet Gynecol 1983;62:539-549.

Waller DK, Keddle AM, Canfield MA, Scheuerle AE. Do infants with major congenital anomalies have an excess of macrosomia? Teratology 2001;64:311-317.

Werler MM, Hayes C, Louik C, Shapiro S, Mitchell AA. Multivitamin supplementation and risk of birth defects. Am J Epidemiol 1999;150:675-682.

Wiswell TE, Tuttle DJ, Northam RS, Simonds GR. Major congenital neurologic malformations. A 17-year survey. Am J Dis Child 1990;144:61-67

Zhang J, Cai WW. Association of the common cold in the first trimester of pregnancy with birth defects. Pediatrics 1993;92:559-563.

Please Note: The primary purpose of this report is to provide background necessary for conducting cluster investigations. It summarizes literature about risk factors associated with this defect. The strengths and limitations of each reference were not critically examined prior to inclusion in this report. Consumers and professionals using this information are advised to consult the references given for more in-depth information.

This report is for information purposes only and is not intended to diagnose, cure, mitigate, treat, or prevent disease or other conditions and is not intended to provide a determination or assessment of the state of health. Individuals affected by this condition should consult their physician and when appropriate, seek genetic counseling.